






Reyaksyon / sentòm apre vaksen an		Kijan sentòm sa a komen?	Kisa pou w fe
Sèlman kote yo te ba w piki an (sètadi, li wouj oswa anfle kote yo te ba w piki an)		<b>Sa rive souvan anpil</b>	Mete konprès frèt kote yo te ba w piki a. Ou ka pran asetaminofèn oswa ibuprofèn pou doulè.
Lejè apre reyaksyon vaksen an, tankou bagay sa yo: <ul style="list-style-type: none"> <li>• Tankou yon fyèv men tanperati ou mwens pase 100.4 degre farennay</li> <li>• Yon ti tèt fè mal</li> <li>• Nouvo fatig (yon fatig ou pa te santi avan)</li> <li>• Yon ti myalji (doulè nan mis)</li> <li>• Yon ti atyalji (doulè atikilasyon)</li> </ul>		<b>Sa rive souvan anpil</b>	Ou ka pran asetaminofèn oswa ibuprofèn si ou gen doulè epi ou santi ou gen lafyèv
Ti sentòm alèji (gratèl, demanjezon men PA itikè) ki rive after apre ou kite sit la		<b>Sa pa rive souvan</b>	Mete konprès frèt kote yo te ba w piki a. Ou ka pran medikaman kont gratèl tankou Claritin, Zyrtec oswa Benadryl.
Sentòm ki modere ak grav, tankou sa yo: <ul style="list-style-type: none"> <li>• Fyèv ki rive 100.4 degre Farennay oswa plis</li> <li>• Tèt fè mal mwayen / grav [oswa]</li> <li>• Fatig mwayen a grav (santi ou fatige) [oswa]</li> <li>• Miyalji (migrèn) mwayen/grav [oswa]</li> <li>• Atyalji mwayen/grav [oswa]</li> </ul>		<b>Sa pa rive souvan</b>	Pran acetaminofèn oswa ibuprofèn pou doulè oswa lafyèv ; rele klinik founisè swen sante ou pou jwenn plis detay.
Reyaksyon alèjik grav tankou itikè, po bouch, lang, je, oswa figi anfle, ral, doulè nan pwtrain oswa souf kout		<b>Ra anpil</b>	Si ou gen difikilte pou respire, bouch oswa gòj ou anfle, oswa ral, rele 911 oswa ale nan sal dijans ki pi pre w la